
Guidelines for Using Point-of-Care SARS-CoV-2 Antigen Tests in Long-Term Care Facilities

Background

Point of care (POC) antigen testing is an important tool to identify and mitigate transmission of SARS-CoV-2 in the long-term care setting. This guidance addresses appropriate use of the tests, understanding results, CLIA requirements, and federal and state reporting mechanisms.

What is an antigen test and how does it differ from the PCR test?

POC antigen tests detect antigens (proteins) on the surface of the SARS-CoV-2 virus while PCR tests detect viral RNA. Given the different detection methods of each, POC antigen tests may not detect the virus as soon as a PCR test. Consequently, false negative results can occur, particularly in the early days of infection. False positive results are rare, particularly with symptomatic individuals. Please see the CDC's [antigen test algorithm](#) to determine when a confirmatory PCR test is needed after a positive or negative POC antigen test result.

When to use the antigen test

The POC antigen test is most useful for testing symptomatic individuals. Since test results can be returned in as little as 15 minutes, a positive result can be used by the facility to quickly decide the type of care necessary for the patient or healthcare worker to prevent transmission to others in the facility. In the context of an outbreak in the long-term care setting, it is recommended to perform COVID-19 outbreak testing every three days if using only POC antigen testing.

While it is generally not recommended to test individuals in the 90 days after they have had a known infection, antigen tests can be used to determine readiness of healthcare workers to return to work after an infection. Please see [CDCs Return to Work Criteria](#) for more information.

What are the CLIA requirements for using the test?

Any facility that performs laboratory tests on human specimens for the purpose of diagnosis and/or treatment is required by federal law to have a Clinical Laboratory Improvement Amendments (CLIA) certificate. There are different types of CLIA certificates; a CLIA Certificate of Waiver covers all tests that are classified as waived by the FDA. Most POC antigen tests are included on the waived list and can be found [here](#) with a 'w' in the authorized settings column. Many facilities have obtained a certificate of waiver previously. For those that have not, instructions and an application for the certificate of waiver can be found [here](#). A CLIA waiver is valid for two years and facilities must renew their certificate of waiver prior to expiration. CLIA certificates can be found using [this search tool](#).

How to collect a specimen safely

Most antigen tests require a swab of the anterior nares. Please follow the manufacturer's instructions for proper specimen collection. Following the manufacturer's instructions will improve reliability of results and is required under the CLIA Certificate of Waiver. Recommended PPE for specimen collection includes an N95, face shield, gown and gloves. Ideally, specimen collection should occur in a private, well-ventilated area.

How to report results to the Utah Department of Health

All COVID-19 test results (positive, negative, equivocal, indeterminate) for all test types (PCR, antigen, serology) are reportable to the Utah Department of Health (UDOH).

- Test results must be reported to UDOH within 24 hours of testing
- Facilities enrolled in the NHSN network, should report all antigen testing results via NHSN
- Please direct POC antigen test result reporting questions to edx@utah.gov

References

Centers for Disease Control and Prevention. (Updated March 4, 2022). Guidance for Antigen Testing for SARS-CoV-2 for Healthcare Providers Testing Individuals in the Community. Retrieved March 28, 2022, from <https://www.cdc.gov/coronavirus/2019-ncov/lab/resources/antigen-tests-guidelines.html>

Centers for Disease Control and Prevention. (Updated Jan. 21, 2022). *Interim guidance for managing healthcare personnel with SARS-COV-2 infection or exposure to SARS-COV-2*. Retrieved March 25, 2022, from https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-risk-assesment-hcp.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fhcp%2Freturn-to-work.html

Centers for Disease Control and Prevention. (Updated Feb. 17, 2022). SARS-COV-2 antigen testing in Long Term Care Facilities. Retrieved March 28, 2022, from <https://www.cdc.gov/coronavirus/2019-ncov/hcp/nursing-homes-antigen-testing.html>

Centers for Medicare and Medicaid Services. (March 2019) *How to Obtain a CLIA Certificate of Waiver*. Retrieved March 28, 2022, from <https://www.cms.gov/regulations-and-guidance/legislation/clia/downloads/howobtaincertificateofwaiver.pdf>